

4. Ramani S, Krackov SK. Twelve tips for giving feedback effectively in the clinical environment. *Med Teach* 2012; 34: 787–91
5. van der Vleuten CPM, Schuwirth LWT, Driessen EW, et al. A model for programmatic assessment fit for purpose. *Med Teach* 2012; 34: 205–14
6. Jivet I, Scheffel M, Schmitz M, et al. From students with love: an empirical study on learner goals, self-regulated learning and sense-making of learning analytics in higher education. *Internet High Educ* 2020; 47: 100758
7. Garrison DR. Self-directed learning: towards a comprehensive model. *Adult Educ Q* 1997; 48: 18–33
8. Hall AK, Nousiainen MT, Campisi P, et al. Training disrupted: practical tips for supporting competency-based medical education during the COVID-19 pandemic. *Med Teach* 2020; 42: 756–61
9. Boscardin C, Fergus KB, Hellevig B, Hauer KE. Twelve tips to promote successful development of a learner performance dashboard within a medical education program. *Med Teach* 2018; 40: 855–61
10. Pingree EW, Huth K, Harper BD, et al. Encouraging entrustment: a qualitative study of resident behaviors that promote entrustment. *Acad Med* 2020. <https://doi.org/10.1097/ACM.0000000000003487> [Epub May 5, 2020 ahead of print]

doi: 10.1016/j.bja.2020.08.016

Advance Access Publication Date: 20 August 2020

© 2020 British Journal of Anaesthesia. Published by Elsevier Ltd. All rights reserved.

## Maintaining education and professional development for anaesthesia trainees during the COVID-19 pandemic: the Self-isolating Virtual Education (SAVED) project

Danielle V. Eusuf<sup>1,\*</sup>, Emma L. England<sup>1</sup>, Mike Charlesworth<sup>2</sup>, Clifford L. Shelton<sup>2,3</sup> and Sarah J. Thornton<sup>4</sup>

<sup>1</sup>North West School of Anaesthesia, Health Education North West, UK, <sup>2</sup>Wythenshawe Hospital, Manchester University NHS Foundation Trust, Manchester, UK, <sup>3</sup>Lancaster Medical School, Faculty of Health and Medicine, Lancaster University, Lancaster, UK and <sup>4</sup>Royal Bolton Hospital, Bolton NHS Foundation Trust, Bolton, UK

\*Corresponding author. E-mail: [danikirk@doctors.org.uk](mailto:danikirk@doctors.org.uk)


**Keywords:** COVID-19; distance learning; educational technology; graduate medical education; self-isolating; social media

Editor—In early 2020, anaesthetists across the world adopted frontline roles in the fight against coronavirus disease 2019 (COVID-19).<sup>1</sup> However, those who were deemed vulnerable to the complications of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection were asked to 'shield' and so self-isolated at home to avoid exposure.<sup>2</sup> Likewise, those who tested positive for SARS-CoV-2 or whose household members had experienced symptoms were asked to self-isolate. Many of these colleagues began to work from home, a challenge considering the 'hands-on' nature of anaesthesia.<sup>3</sup> Motivated by a wish to make a meaningful contribution during the pandemic, and in response to the ongoing educational need despite the postponement of face-to-face tutorials, the Self-isolating Virtual Education (SAVED) project was created by self-isolating trainees based in the North West School of Anaesthesia, UK. In this report, we describe the design of this online educational intervention, and present its impact in terms of uptake and educational evaluation. The Health Education England Research Governance Group granted permission to collect and present this data (date of approval: May 19, 2020).

Based on the Royal College of Anaesthetists' training curriculum, with a focus on preparation for examinations, self-isolating trainees designed and hosted a combination of pre-recorded and live online tutorials. These were delivered

using a combination of videoconferencing (Zoom Version 5.0.2; Zoom Video Communications Inc., San Jose, CA, USA) and video-sharing (Vimeo Pro; Vimeo Inc., New York, NY, USA). At the time of writing, 24 live tutorials and two live study days had been delivered, and more than 80 pre-recorded tutorials had been made available through the North West School of Anaesthesia website ([www.mmacc.uk](http://www.mmacc.uk)). The tutorials were predominantly delivered by trainees, with some taught by consultants; topics were appropriately matched to clinical experience and expertise. All recorded tutorials were peer-reviewed by a senior trainee or a consultant before being published online. The resources were promoted through existing trainee e-mail contact lists, social media, and by sharing resources with other deaneries.

We were able to quantify the reach of SAVED using social media platform analytics and website metrics.<sup>4</sup> Between March 27 and June 27, 2020, we recorded 4881 visits to the [www.mmacc.uk](http://www.mmacc.uk) website; the video tutorials received 8304 impressions (number of times the video web page was visited) and 3720 views (number of times the video was played) (Fig. 1). Content was accessed from 50 cities within the UK and 27 countries outside the UK. By June 27, 2020, the @SAVED\_anaes Twitter account, which was established on April 8, recorded ~54 000 impressions, 1528 profile visits, 270 followers, 245 retweets, 382 likes, 763 link clicks, and a 6.6% engagement rate.



**Fig 1.** Daily impressions (light blue) and views (dark blue) of SAVED video resources on Vimeo, and the timing of promotional interventions between March 27 and June 27, 2020. E, e-mails; S, social media accounts opened; D, resources shared with other deaneries; W, national webinar.

The use of e-mail and social media appeared to have an important impact in stimulating engagement with the project. There has been an increasing trend in the number of impressions and views of the tutorials over time with 'spikes' after interventions such as reminder e-mails, the opening of Twitter and Facebook accounts, and the participation of one of the SAVED faculty as a panellist in a national webinar.<sup>5</sup> Likewise, an increase in engagement was seen after sharing of the resources with other deaneries (Fig 1).

We assessed the educational impact of the SAVED project using post-tutorial feedback and, at the preference of the presenter in some live tutorials ( $n=3$ ), pre- and post-tutorial multiple choice questions (MCQs) administered using the polling function. Attendees rated the resources using 5-point Likert scales and optionally, free-text feedback. We received 456 completed feedback forms which showed median (inter-quartile range) scores for quality of content, presentation of material and usefulness compared with traditional tutorials of 5 (5–5), 5 (4–5) and 5 (4–5), respectively. MCQ scores were higher post-tutorial than pre-tutorial in the three tutorials where they were used (mean improvement of 24%, 22%, and 32%). Free-text feedback highlighted benefits including less time spent travelling and the ability to access learning more flexibly. Disadvantages included the loss of social contact between trainees that might normally occur during the break-times of face-to-face tutorials,<sup>6</sup> and less interaction between attendees and facilitators. Social isolation has a role in physician burnout,<sup>7</sup> a relevant problem during the COVID-19 pandemic. We therefore attempted to facilitate social

support by continuing the session after the facilitator left so that attendees were able to engage in informal conversation. The addition of live MCQ polls and the use of pre-tutorial materials notably enhanced attendee–facilitator interaction. Anecdotally, interaction has also improved over time as users have become more at ease with the technology.

Trainees were allocated time away from clinical work to attend live tutorials; therefore, attendance was expected. Nevertheless, more people attended live virtual sessions than would normally be expected for face-to-face tutorials. These were also found to be more reliable as there were fewer cancellations by facilitators owing to clinical commitments. Accessing the pre-recorded online tutorials was entirely optional and therefore dependant on trainee self-direction. Although a small incentive in the form of continuing professional development (CPD) points was available for attending tutorials, this is unlikely to have had a major impact on uptake; previous work has found that the motivation to engage with online learning is predominantly influenced not by external factors but by interest, control, and freedom of choice,<sup>8</sup> all of which are features of the approach we adopted.

The SAVED project has given self-isolating trainees the ability to work from home to create online tutorials that have been well received and widely utilised by colleagues in the North West School of Anaesthesia and beyond. Social media and online sharing have been instrumental in promoting the uptake of these resources. Although we are keen that face-to-face tutorials should restart when possible, the success of this project has provided a new vision for anaesthetic training in

the North West. We envisage an increasing role for online education to supplement traditional teaching allowing for greater flexibility and hopefully increased engagement. In this new era of free access to online medical resources and technology-enhanced learning,<sup>9,10</sup> we are confident that our resource and others like it will continue to grow and make a substantial contribution to anaesthetic training in the UK and around the world.

## Acknowledgements

The authors acknowledge the contribution of all those who produced content for the SAVED project during the COVID-19 pandemic, and John-Paul Lomas for his assistance in providing website usage data.

## Declarations of interest

CS is a former member of the editorial board of *BJA Education*. The other authors declare no other competing interests.

## References

- Ortega R, Chen R. Beyond the operating room: the roles of anaesthesiologists in pandemics. *Br J Anaesth* 2020. <https://doi.org/10.1016/j.bja.2020.06.005>. Advance Access published on June 11, 2020
- Illiff HA, Simpson KA, Tomlinson CR, Webb CM. 'Shielded' anaesthetists and intensivists during the COVID-19 pandemic. *Anaesthesia* 2020. <https://doi.org/10.1111/anae.15153>. Advance Access published on July 6
- World Health Organization. *Mental health and psychosocial considerations during the COVID-19 outbreak* March 2020. Available from: <https://apps.who.int/iris/bitstream/handle/10665/331490/WHO-2019-nCoV-MentalHealth-2020.1-eng.pdf>. [Accessed 10 June 2020]
- Johannsson H, Selak T. Dissemination of medical publications on social media – is it the new standard? *Anaesthesia* 2020; 75: 155–7
- Intensive Care Society. *Education in COVID* June 2020. Available from: <https://youtu.be/elwjDG97GZw>. [Accessed 2 July 2020]
- Muilenburg LY, Berge ZL. Student barriers to online learning: a factor analytic study. *Distance Educ* 2005; 26: 29–48
- Wong AVK, Olusanya O. Burnout and resilience in anaesthesia and intensive care medicine. *BJA Educ* 2017; 17: 334–40
- Song D, Bonk CJ. Motivational factors in self-directed informal learning from online learning resources. *Cogent Educ* 2016; 3: 1–11
- Carroll CL, Bruno K, vonTschudi M. Social media and free open access medical education: the future of medical and nursing education? *Am J Crit Care* 2016; 25: 93–6
- Wang JJ, Deng A, Tsui BCH. COVID-19: novel pandemic, novel generation of medical students. *Br J Anaesth* 2020. <https://doi.org/10.1016/j.bja.2020.06.005>. Advance Access published on June 2

doi: 10.1016/j.bja.2020.07.046

Advance Access Publication Date: 7 August 2020

© 2020 British Journal of Anaesthesia. Published by Elsevier Ltd. All rights reserved.

# Bedside monitoring of lung perfusion by electrical impedance tomography in the time of COVID-19

Bijan Safaee Fakhr<sup>1,2</sup>, Caio C. Araujo Morais<sup>1,2</sup>, Roberta R. De Santis Santiago<sup>1,2</sup>, Raffaele Di Fenza<sup>1,2</sup>, Lauren E. Gibson<sup>1,2</sup>, Paula A. Restrepo<sup>1,2</sup>, Marvin G. Chang<sup>1,2</sup>, Edward A. Bittner<sup>1,2</sup>, Riccardo Pinciroli<sup>1,2</sup>, Florian J. Fintelmann<sup>2,3</sup>, Robert M. Kacmarek<sup>1,2,4</sup> and Lorenzo Berra<sup>1,2,4,\*</sup>

<sup>1</sup>Department of Anaesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital, Boston, MA, USA, <sup>2</sup>Harvard Medical School, Boston, MA, USA, <sup>3</sup>Department of Radiology, Massachusetts General Hospital, Boston, MA, USA and <sup>4</sup>Respiratory Care Department, Massachusetts General Hospital, Boston, MA, USA

\*Corresponding author. E-mail: [lberra@mgh.harvard.edu](mailto:lberra@mgh.harvard.edu)

**Keywords:** COVID-19; electrical impedance tomography; lung perfusion; mechanical ventilation; pulmonary embolism; ventilation/perfusion matching

Editor—Coronavirus disease 2019 (COVID-19) results in a broad spectrum of clinical presentations, including viral pneumonia and acute respiratory distress syndrome (ARDS).<sup>1</sup> Increasingly compelling evidence suggests that the underlying pathophysiology of severe COVID-19 pneumonia is microvascular thrombosis.<sup>2,3</sup> Although the pulmonary

vasculature can be seen in exquisite detail with computed tomography pulmonary angiography (CTPA), routine CTPA is neither feasible nor recommended in COVID-19 patients.<sup>4</sup> Therefore, tools enabling bedside evaluation of lung perfusion, including monitoring the response to therapeutic